



For adults with follicular lymphoma that has relapsed (*went into remission, then came back*) or is refractory (*did not go into remission after receiving other lymphoma treatments*) after having at least 2 other kinds of treatment

BREAK FREE FROM THE CYCLE

A SINGLE INFUSION. A CHANCE AT LASTING REMISSION.

What is KYMRIA[®] (tisagenlecleucel)?

KYMRIA is made from your own white blood cells and is a prescription cancer treatment used in patients with follicular lymphoma (FL), a type of non-Hodgkin lymphoma, that has relapsed (went into remission, then came back) or is refractory (did not go into remission after receiving other lymphoma treatments) after having at least two other kinds of treatment.

There are ongoing studies to confirm the benefit of KYMRIA for FL.

IMPORTANT SAFETY INFORMATION

What is the most important information I should know about KYMRIA?

KYMRIA may cause side effects that are severe or life-threatening, such as cytokine release syndrome (CRS) and neurological toxicities. Call your health care provider or get emergency help right away if you get any of the following signs and symptoms of Cytokine Release Syndrome:

Please see additional Important Safety Information throughout and Summary of Important Information on pages 13-15.



KYMRIAH OFFERS A CHANCE AT LASTING REMISSION



KYMRIAH enhances the ability of your T cells to detect and destroy your healthy and cancerous B cells



KYMRIAH is typically given in a single infusion made up of your own immune cells



Collaboration with your treatment team and caregivers is imperative

Making treatment decisions for relapsed or refractory follicular lymphoma can be challenging. Talk with your treatment team about questions you may have.

Visit KYMRIAH.com to learn more about KYMRIAH, treatment centers, and available support

IMPORTANT SAFETY INFORMATION (cont)

What is the most important information I should know about KYMRIAH? (cont)

• Cytokine Release Syndrome:

- Difficulty breathing
- Fever (100.4°F/38°C or higher)
- Chills/shaking chills
- Severe nausea, vomiting, diarrhea
- Severe muscle or joint pain
- Very low blood pressure
- Dizziness/lightheadedness

Please see additional Important Safety Information throughout and Summary of Important Information on pages 13-15.

WHAT IS RELAPSED/REFRACTORY FOLLICULAR LYMPHOMA?

The immune system is made up of different cells and organs that work together to protect the body from diseases. For example, there are B cells, T cells, and glands called lymph nodes. Sometimes the cells inside a lymph node can grow abnormally and become cancerous.

Follicular lymphoma is a type of non-Hodgkin lymphoma (cancer of the immune system) that develops from B cells and is usually slow growing.

When your cancer is in **remission**, it means the signs and symptoms of cancer have decreased or disappeared. Your cancer is considered to have **relapsed** when it comes back after a period of remission, and to be **refractory** when it does not respond to treatment.


IMPORTANT SAFETY INFORMATION (cont)

What is the most important information I should know about KYMRIAH? (cont)

• Neurological Toxicities:

- Altered or decreased consciousness
- Delirium
- Confusion
- Agitation
- Seizures
- Difficulty speaking and understanding
- Loss of balance


Please see additional Important Safety Information throughout and Summary of Important Information on pages 13-15.



T cells
T cells are a key part of your immune system that can find and destroy cells that are infected or that have become cancerous.



B cells
B cells are also a key component of your immune system. They make special proteins called antibodies, which attach to the surface of foreign invaders, alerting your body to the presence of intruders. Some B cells can then remember the intruders so if they see them again in the future, they can respond more quickly.



Cancerous cells
Your healthy B cells can sometimes change and develop into a cancer, including DLBCL. These abnormal B cells continue to grow and increase in number, which can affect your body in different ways.



KYMRIAH IS A SINGLE INFUSION MADE FROM YOUR OWN IMMUNE CELLS

Unlike traditional chemotherapy or stem cell transplant, KYMRIAH is a type of immunotherapy, called CAR-T cell therapy or chimeric antigen receptor T cell therapy. KYMRIAH is made up of your own T cells that have been reprogrammed to recognize the antigen CD19, allowing them to better detect and destroy cancerous (and normal) B cells in the body.

KYMRIAH is approved in patients with follicular lymphoma that has relapsed (went into remission, then came back) or is refractory (did not go into remission after receiving other lymphoma treatments) after having at least 2 other kinds of treatment.


KYMRIAH is a **single infusion** that may allow patients to break free from the cycle of daily oral medications and multiple intravenous infusions.

IMPORTANT SAFETY INFORMATION (cont)


What is the most important information I should know about KYMRIAH? (cont)

You may be admitted to the hospital and treated with other medications if you have any of these side effects. If you are admitted to the hospital, tell the health care provider that you have received KYMRIAH.

Please see additional Important Safety Information throughout and Summary of Important Information on pages 13-15.



WHAT IS THE PROCESS FOR GETTING KYMRIAH?*



*KYMRIAH is only available at select treatment centers.

Please see additional Important Safety Information throughout and Summary of Important Information on pages 13-15.



KYMRIAH IS A POTENTIALLY DEFINITIVE, SINGLE INFUSION THAT MAY BREAK THE CYCLE OF REPEATED TREATMENTS IN ADULT PATIENTS WITH RELAPSED OR REFRACTORY FOLLICULAR LYMPHOMA


In a global, phase 2 pivotal study (ELARA), **82%** of patients did not require another anti-lymphoma treatment within 12 months post infusion



Multiple IV Infusions



Daily Oral Medications



Indefinite Repeated Treatment Cycles

IMPORTANT SAFETY INFORMATION (cont)


What are other serious side effects of KYMRIAH?

- **Hemophagocytic Lymphohistiocytosis (HLH)/Macrophage Activation Syndrome (MAS):** Be sure to discuss with your health care provider the possibility of developing this life-threatening condition, and thereafter, your doctor will monitor you for the possibility of developing HLH/MAS
- **Allergic Reactions:** Serious allergic reactions, including anaphylaxis, which is a life-threatening allergic reaction, may occur after you receive KYMRIAH. Some signs and symptoms may include difficulty breathing, very low blood pressure, dizziness, swelling under skin, rash, nausea, and vomiting. You should seek emergency medical treatment right away if you have an allergic reaction
- **Serious Infections:** KYMRIAH can increase the risk of life-threatening infections that may lead to death. Tell your health care provider right away if you develop fever, chills, or any signs or symptoms of an infection

Please see additional Important Safety Information throughout and Summary of Important Information on pages 13-15.

NEARLY 90% OF PATIENTS RESPONDED TO TREATMENT WITH KYMRIAH

In the phase 2 ELARA clinical trial, KYMRIAH delivered strong efficacy in adult patients



- 86%** (77 of 90) of patients **responded to treatment with KYMRIAH**
- 68%** (61 of 90) of patients achieved a **complete response**
 - All signs of cancer disappeared in the body in 68% of adults
- 18%** (16 of 90) of patients achieved a **partial response**
 - 18% of adults have shown a decrease in cancer cells within the body

IMPORTANT SAFETY INFORMATION (cont)

What are other serious side effects of KYMRIAH? (cont)

- **Prolonged Low Blood Cell Counts (Cytopenias):** KYMRIAH can lower 1 or more types of your blood cells (red blood cells, white blood cells, or platelets). After treatment, your health care provider may test your blood to check cell counts. Tell your health care provider right away if you get a fever or other symptoms of an infection, are feeling tired, weak, or short of breath, or have unusual bruising or bleeding
- **Hypogammaglobulinemia:** A condition in which the level of immunoglobulins (antibodies) in your blood is low and the risk of infection is increased. It is expected that you may develop hypogammaglobulinemia with KYMRIAH, and you may need to receive immunoglobulin replacement for an indefinite amount of time following treatment with KYMRIAH. Tell your health care provider about your treatment with KYMRIAH before receiving a live vaccine

Please see additional Important Safety Information throughout and Summary of Important Information on pages 13-15.



WHAT SIDE EFFECTS MAY I EXPERIENCE?

KYMRIAH® (tisagenlecleucel) may cause side effects that are severe or life-threatening. Your treatment team is specifically trained to monitor for and manage these potential side effects. Most side effects happen in the weeks following infusion with KYMRIAH.

Call your health care provider or seek emergency help right away if you experience any of the following:

 Difficulty breathing	 Severe muscle or joint pain	 Loss of balance
 Fever (100.4 °F/38 °C or higher)	 Very low blood pressure	 Seizures
 Chills/shaking chills	 Dizziness/lightheadedness	 Difficulty speaking and understanding
 Confusion		

You may be admitted to the hospital and treated with other medications if you have any of these side effects. If you are admitted to the hospital, tell the health care provider that you have received KYMRIAH.

These are not all the possible side effects of KYMRIAH. Please see additional important safety information throughout and Summary of Important Safety Information on pages 13-15

IMPORTANT SAFETY INFORMATION (cont)

What are other serious side effects of KYMRIAH? (cont)

- **Secondary Cancers:** KYMRIAH may increase your risk of getting cancers, including certain types of blood cancers. Your health care provider should monitor you for this

IMPORTANT SAFETY INFORMATION (cont)

How will I get KYMRIAH?

- Since KYMRIAH is made from your own white blood cells, your health care provider has to take some of your blood. This is called “leukapheresis.” It takes 3 to 6 hours and may need to be repeated. A tube (intravenous catheter) will be placed in your vein to collect your blood
- Your blood cells are frozen and sent to the manufacturing site to make KYMRIAH. It takes about 3 to 4 weeks from the time your cells are received at the manufacturing site and shipped to your health care provider, but the time may vary
- While waiting for KYMRIAH to be made, your health care provider may give you therapy to stabilize your cancer
- In addition, before you get KYMRIAH, your health care provider may give you chemotherapy for a few days to prepare your body. When your body is ready, your health care provider will give you KYMRIAH through a tube (intravenous catheter) in your vein. This usually takes less than 1 hour
- You should plan to stay close to a health care facility for at least 2 weeks after getting KYMRIAH. Your health care provider will check to see if your treatment is working and help you with any side effects that occur

WHAT SHOULD I ASK MY TREATMENT TEAM?

Knowing who is on your treatment team for KYMRIA[®]H and how to get in touch with them are important. Remember to ask for names, telephone numbers, email addresses, and any other contact information.

Initial questions might include the following:

- How does my treatment history affect my eligibility for KYMRIA[®]H?
- Can I receive KYMRIA[®]H after a bone marrow transplant?
- What if I have other health conditions?
- How will I feel throughout the treatment process?
- Where can I receive KYMRIA[®]H therapy?
- How long will I need to stay near the treatment center?
- Will my insurance cover KYMRIA[®]H therapy?

Click [here](#) for a discussion guide to use with your doctor.



IMPORTANT SAFETY INFORMATION (cont)

What are the possible or reasonably likely side effects of KYMRIA[®]H?

The most common side effects of KYMRIA[®]H include:

- | | |
|-------------------------------------|-------------------------------|
| • Difficulty breathing | • Severe muscle or joint pain |
| • Fever (100.4°F/38°C or higher) | • Very low blood pressure |
| • Chills/shaking chills | • Dizziness/lightheadedness |
| • Confusion | • Headache |
| • Severe nausea, vomiting, diarrhea | |

These are not all the possible side effects of KYMRIA[®]H. Talk to your health care provider for medical advice about side effects.

Please see additional Important Safety Information throughout and Summary of Important Information on pages 13-15.

WHAT SHOULD I ASK MY TREATMENT TEAM?

Collection

- How should I prepare for the collection?
- How long will it take?
- What can I do during collection?
- Will there be any limitations or side effects afterward?
- How soon after collection will my KYMRIA[®]H CAR-T cells be ready for infusion?

Preinfusion Lymphodepleting Chemotherapy

- How is this chemotherapy different from the chemotherapy I previously received?
- How long will it take?
- Will there be side effects?
- How far in advance of my infusion do I receive chemotherapy?

Infusion

- How should I prepare for infusion?
- What will infusion be like?
- How long will it take?

Side Effects and Monitoring

- What side effects should I expect after infusion?
- How will serious side effects be managed after therapy?
- How will I know KYMRIA[®]H is working?
- When do I need to check in with my treatment team?
- How long do I need to stay in or near my hospital?
- After returning home, what kind of side effects require a hospital visit?
- How quickly can I get back to my daily routine?

IMPORTANT SAFETY INFORMATION (cont)

What should I tell my health care provider before receiving KYMRIA[®]H?

- Tell your health care provider if you are pregnant, planning to be pregnant, or breastfeeding. Your health care provider may do a pregnancy test prior to your starting treatment. No information is available of KYMRIA[®]H use in pregnant or breastfeeding women. Therefore, KYMRIA[®]H is not recommended for women who are pregnant or breastfeeding. Talk to your health care provider about birth control and pregnancy.
- Tell your health care provider about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements.

Please see additional Important Safety Information throughout and Summary of Important Information on pages 13-15.



AS A CAREGIVER, HOW I CAN HELP THROUGHOUT THE PROCESS?

Finding out your loved one’s cancer has returned or hasn’t responded to treatment can be hard on everyone’s mental and emotional well-being. It is important to tap into available resources and practice self-care during this difficult time. Leaning on social and emotional support networks, engaging in physical activity, and connecting with spirituality can be helpful.

You can help by watching for side effects. KYMRIA[®]H can be given in an inpatient (stay in the center) or outpatient (leave center after infusion) setting. Before leaving the treatment center, your treatment team will give you a list of potential side effects to watch for, as well as instructions for what to do if they happen.

KYMRIA[®]H is only available at select treatment centers. You should plan to stay close to a health care facility for at least 2 weeks after getting KYMRIA[®]H. Your health care provider will check to see if your treatment is working and help you with any side effects that occur.



IMPORTANT SAFETY INFORMATION (cont)

What should I be aware of after receiving KYMRIA[®]H?

- Avoid driving for at least 2 weeks after you get KYMRIA[®]H
- Do not donate blood, organs, tissues, sperm, oocytes, and other cells
- Some commercial HIV tests may cause a false positive HIV test result

This is a summary of the most important safety information about KYMRIA[®]H. Talk with your health care provider or pharmacist about side effects. If you would like more information, the FDA-approved product labeling for KYMRIA[®]H can be found at www.KYMRIA.com or call 1-888-669-6682.

You are encouraged to report negative side effects of prescription drugs to the FDA. Visit www.fda.gov/medwatch or call 1-800-FDA-1088.

Please see additional Important Safety Information throughout and Summary of Important Information on pages 13-15.

SUMMARY OF IMPORTANT SAFETY INFORMATION

What is KYMRIA[®]H?

KYMRIA[®]H is made from your own white blood cells and is a prescription cancer treatment used in patients with follicular lymphoma (FL), a type of non-Hodgkin lymphoma, that has relapsed (went into remission, then came back) or is refractory (did not go into remission after receiving other lymphoma treatments) after having at least two other kinds of treatment.

There are ongoing studies to confirm the benefit of KYMRIA[®]H in FL.

What is the most important information I should know about KYMRIA[®]H?

KYMRIA[®]H may cause side effects that are severe or life-threatening, such as cytokine release syndrome (CRS) and neurological toxicities. Call your health care provider or get emergency help right away if you get any of the following signs and symptoms of:

• Cytokine Release Syndrome:

- Difficulty breathing
- Fever (100.4°F/38°C or higher)
- Chills/shaking chills
- Severe nausea, vomiting, diarrhea
- Severe muscle or joint pain
- Very low blood pressure
- Dizziness/lightheadedness

• Neurological Toxicities:

- Altered or decreased consciousness
- Delirium
- Confusion
- Agitation
- Seizures
- Difficulty speaking and understanding
- Loss of balance

You may be admitted to the hospital and treated with other medications if you have any of these side effects. If you are admitted to the hospital, tell the health care provider that you have received KYMRIA[®]H.

Please see additional Important Safety Information throughout and Summary of Important Information on pages 13-15.



SUMMARY OF IMPORTANT SAFETY INFORMATION (continued)

What are other serious side effects of KYMRIA?

- **Hemophagocytic Lymphohistiocytosis (HLH)/Macrophage Activation Syndrome (MAS):** Be sure to discuss with your health care provider the possibility of developing this life-threatening condition, and thereafter, your doctor will monitor you for the possibility of developing HLH/MAS
- **Allergic Reactions:** Serious allergic reactions, including anaphylaxis, which is a life-threatening allergic reaction, may occur after you receive KYMRIA. Some signs and symptoms may include difficulty breathing, very low blood pressure, dizziness, swelling under skin, rash, nausea, and vomiting. You should seek emergency medical treatment right away if you have an allergic reaction
- **Serious Infections:** KYMRIA can increase the risk of life-threatening infections that may lead to death. Tell your health care provider right away if you develop fever, chills, or any signs or symptoms of an infection
- **Prolonged Low Blood Cell Counts (Cytopenias):** KYMRIA can lower 1 or more types of your blood cells (red blood cells, white blood cells, or platelets). After treatment, your health care provider may test your blood to check cell counts. Tell your health care provider right away if you get a fever or other symptoms of an infection, are feeling tired, weak, or short of breath, or have unusual bruising or bleeding
- **Hypogammaglobulinemia:** A condition in which the level of immunoglobulins (antibodies) in your blood is low and the risk of infection is increased. It is expected that you may develop hypogammaglobulinemia with KYMRIA, and you may need to receive immunoglobulin replacement for an indefinite amount of time following treatment with KYMRIA. Tell your health care provider about your treatment with KYMRIA before receiving a live vaccine
- **Secondary Cancers:** KYMRIA may increase your risk of getting cancers, including certain types of blood cancers. Your health care provider should monitor you for this

How will I get KYMRIA?

- Since KYMRIA is made from your own white blood cells, your health care provider has to take some of your blood. This is called “leukapheresis.” It takes 3 to 6 hours and may need to be repeated. A tube (intravenous catheter) will be placed in your vein to collect your blood
- Your blood cells are frozen and sent to the manufacturing site to make KYMRIA. It takes about 3 to 4 weeks from the time your cells are received at the manufacturing site and shipped to your health care provider, but the time may vary
- While waiting for KYMRIA to be made, your health care provider may give you therapy to stabilize your cancer

SUMMARY OF IMPORTANT SAFETY INFORMATION (continued)

- In addition, before you get KYMRIA, your health care provider may give you chemotherapy for a few days to prepare your body. When your body is ready, your health care provider will give you KYMRIA through a tube (intravenous catheter) in your vein. This usually takes less than 1 hour
- You should plan to stay close to a health care facility for at least 2 weeks after getting KYMRIA. Your health care provider will check to see if your treatment is working and help you with any side effects that occur

What are the possible or reasonably likely side effects of KYMRIA?

The most common side effects of KYMRIA include:

- | | | |
|----------------------------------|-------------------------------------|-----------------------------|
| • Difficulty breathing | • Confusion | • Very low blood pressure |
| • Fever (100.4°F/38°C or higher) | • Severe nausea, vomiting, diarrhea | • Dizziness/lightheadedness |
| • Chills/shaking chills | • Severe muscle or joint pain | • Headache |

These are not all the possible side effects of KYMRIA. Talk to your health care provider for medical advice about side effects.

What should I tell my health care provider before receiving KYMRIA?

- Tell your health care provider if you are pregnant, planning to be pregnant, or breastfeeding. Your health care provider may do a pregnancy test prior to your starting treatment. No information is available of KYMRIA use in pregnant or breastfeeding women. Therefore, KYMRIA is not recommended for women who are pregnant or breastfeeding. Talk to your health care provider about birth control and pregnancy
- Tell your health care provider about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements

What should I be aware of after receiving KYMRIA?

- Avoid driving for at least 2 weeks after you get KYMRIA
- Do not donate blood, organs, tissues, sperm, oocytes, and other cells
- Some commercial HIV tests may cause a false positive HIV test result

This is a summary of the most important safety information about KYMRIA. Talk with your health care provider or pharmacist about side effects. If you would like more information, the FDA-approved product labeling for KYMRIA can be found at www.KYMRIA.com, or call 1-888-669-6682.

You are encouraged to report negative side effects of prescription drugs to the FDA. Visit www.fda.gov/medwatch, or call 1-800-FDA-1088.

Please see additional Important Safety Information throughout and Summary of Important Information on pages 13-15.



Please see additional Important Safety Information throughout and Summary of Important Information on pages 13-15.

WHAT KIND OF SUPPORT IS AVAILABLE?

American Cancer Society
www.Cancer.org

Friends of Cancer Research
www.FOCR.org

Patient Advocate Foundation
www.PatientAdvocate.org

CancerCare
www.CancerCare.org

Blood Cancer United
<https://bloodcancerunited.org>

Cancer Support Community
www.CancerSupportCommunity.org

Cancer Research Institute
www.CancerResearch.org

Lymphoma Research Foundation
www.Lymphoma.org

The websites listed on this page are maintained by third parties over whom Novartis Pharmaceuticals Corporation has no control. As such, Novartis Pharmaceuticals Corporation makes no representation as to the accuracy or any other aspect of the information supplied by these organizations or contained on these websites. The organizations listed on this page do not specifically endorse the use of KYMRIA[®].

For information about financial assistance, patient support programs, or finding a treatment center, contact



To learn more, please call KYMRIA CARES[™] at (1-844-459-6742) from 8:00 AM to 8:00 PM ET.

Please see additional Important Safety Information throughout and Summary of Important Information on pages 13-15.

